

2000 AIME II

For more practice and resources, visit ziml.areteem.org

The problems in the AMC-Series Contests are copyrighted by American Mathematics Competitions at Mathematical Association of America (www.maa.org).



| Question 1 Not yet answered Points out of 5 | The number $\frac{2}{\log_4 2000^6} + \frac{3}{\log_5 2000^6}$ can be written as $\frac{m}{n}$ where m and n are relatively prime positive integers. Find $m + n$. |
|--|---|
| Question 2 Not yet answered Points out of 5 | A point whose coordinates are both integers is called a lattice point. How many lattice points lie on the hyperbola $x^2 - y^2 = 2000^2$? Answer: |
| Question 3 Not yet answered Points out of 5 | A deck of forty cards consists of four 1's, four 2's,, and four 10's. A matching pair (two cards with the same number) is removed from the deck. Given that these cards are not returned to the deck, let m/n be the probability that two randomly selected cards also form a pair, where m and n are relatively prime positive integers. Find $m + n$. Answer: |
| Question 4 Not yet answered Points out of 5 | What is the smallest positive integer with six positive odd integer divisors and twelve positive even integer divisors? Answer: |
| Question 5 Not yet answered Points out of 5 | Given eight distinguishable rings, let n be the number of possible five-ring arrangements on the four fingers (not the thumb) of one hand. The order of rings on each finger is significant, but it is not required that each finger have a ring. Find the leftmost three nonzero digits of n . |

| Question 6 Not yet answered Points out of 5 | One base of a trapezoid is 100 units longer than the other base. The segment that joins the midpoints of the legs divides the trapezoid into two regions whose areas are in the ratio $2:3$. Let x be the length of the segment joining the legs of the trapezoid that is parallel to the bases and that divides the trapezoid into two regions of equal area. Find the greatest integer that does not exceed $x^2/100$. |
|---|--|
| Question 7 Not yet answered Points out of 5 | Given that $\frac{1}{2!17!} + \frac{1}{3!16!} + \frac{1}{4!15!} + \frac{1}{5!14!} + \frac{1}{6!13!} + \frac{1}{7!12!} + \frac{1}{8!11!} + \frac{1}{9!10!} = \frac{N}{1!18!}$ find the greatest integer that is less than $\frac{N}{100}$. Answer: |
| Question 8 Not yet answered Points out of 5 | In trapezoid $ABCD$, leg \overline{BC} is perpendicular to bases \overline{AB} and \overline{CD} , and diagonals \overline{AC} and \overline{BD} are perpendicular. Given that $AB = \sqrt{11}$ and $AD = \sqrt{1001}$, find BC^2 . Answer: |
| Question 9 Not yet answered Points out of 5 | Given that z is a complex number such that $z + \frac{1}{z} = 2\cos 3^\circ$, find the least integer that is greater than $z^{2000} + \frac{1}{z^{2000}}$. Answer: |
| Question 10 Not yet answered Points out of 5 | A circle is inscribed in quadrilateral $ABCD$, tangent to \overline{AB} at P and to \overline{CD} at Q . Given that $AP = 19$, $PB = 26$, $CQ = 37$, and $QD = 23$, find the square of the radius of the circle. |

| Question 11 Not yet answered Points out of 5 | The coordinates of the vertices of isosceles trapezoid $ABCD$ are all integers, with $A = (20, 100)$ and $D = (21, 107)$. The trapezoid has no horizontal or vertical sides, and \overline{AB} and \overline{CD} are the only parallel sides. The sum of the absolute values of all possible slopes for \overline{AB} is m/n , where m and n are relatively prime positive integers. Find $m + n$. Answer: |
|---|--|
| Question 12 Not yet answered Points out of 5 | The points A , B and C lie on the surface of a sphere with center O and radius 20. It is given that $AB = 13$, $BC = 14$, $CA = 15$, and that the distance from O to $\triangle ABC$ is $\frac{m\sqrt{n}}{k}$, where m , n , and k are positive integers, m and k are relatively prime, and n is not divisible by the square of any prime. Find $m + n + k$. Answer: |
| Question 13 Not yet answered Points out of 5 | The equation $2000x^6 + 100x^5 + 10x^3 + x - 2 = 0$ has exactly two real roots, one of which is $\frac{m+\sqrt{n}}{r}$, where m , n and r are integers, m and r are relatively prime, and $r > 0$. Find $m + n + r$. |
| Question 14 Not yet answered Points out of 5 | Every positive integer k has a unique factorial base expansion $(f_1, f_2, f_3, \ldots, f_m)$, meaning that $k = 1! \cdot f_1 + 2! \cdot f_2 + 3! \cdot f_3 + \cdots + m! \cdot f_m$, where each f_i is an integer, $0 \le f_i \le i$, and $0 < f_m$. Given that $(f_1, f_2, f_3, \ldots, f_j)$ is the factorial base expansion of $16! - 32! + 48! - 64! + \cdots + 1968! - 1984! + 2000!$, find the value of $f_1 - f_2 + f_3 - f_4 + \cdots + (-1)^{j+1} f_j$. Answer: |
| Question 15 Not yet answered Points out of 5 | Find the least positive integer n such that $\frac{1}{\sin 45^{\circ} \sin 46^{\circ}} + \frac{1}{\sin 47^{\circ} \sin 48^{\circ}} + \dots + \frac{1}{\sin 133^{\circ} \sin 134^{\circ}} = \frac{1}{\sin n^{\circ}}.$ Answer: |