

2011 AIME I

For more practice and resources, visit ziml.areteem.org

The problems in the AMC-Series Contests are copyrighted by American Mathematics Competitions at Mathematical Association of America (www.maa.org).



| Question 1 Not yet answered Points out of 5 | Jar A contains four liters of a solution that is 45% acid. Jar B contains five liters of a solution that is 48% acid. Jar C contains one liter of a solution that is $k\%$ acid. From jar C, $\frac{m}{n}$ liters of the solution is added to jar A, and the remainder of the solution in jar C is added to jar B. At the end both jar A and jar B contain solutions that are 50% acid. Given that m and n are relatively prime positive integers, find $k + m + n$. |
|--|---|
| | |
| Question 2 Not yet answered Points out of 5 | In rectangle $ABCD$, $AB = 12$ and $BC = 10$. Points E and F lie inside rectangle $ABCD$ so that $BE = 9, DF = 8, \overline{BE} \overline{DF}, \overline{EF} \overline{AB}$, and line BE intersects segment \overline{AD} . The length EF can be expressed in the form $m\sqrt{n} - p$, where m,n , and p are positive integers and n is not divisible by the square of any prime. Find $m + n + p$. Answer: |
| | |
| Question 3 Not yet answered Points out of 5 | Let L be the line with slope $\frac{5}{12}$ that contains the point $A = (24, -1)$, and let M be the line perpendicular to line L that contains the point $B = (5, 6)$. The original coordinate axes are erased, and line L is made the x -axis and line M the y -axis. In the new coordinate system, point A is on the positive x -axis, and point B is on the positive y -axis. The point P with coordinates $(-14, 27)$ in the original system has coordinates (α, β) in the new coordinate system. Find $\alpha + \beta$. |
| | |
| Question 4 Not yet answered Points out of 5 | In triangle ABC , $AB = 125$, $AC = 117$ and $BC = 120$. The angle bisector of angle A intersects \overline{BC} at point L , and the angle bisector of angle B intersects \overline{AC} at point K . Let M and N be the feet of the perpendiculars from C to \overline{BK} and \overline{AL} , respectively. Find MN . |
| | Answer: |

| Question 5 Not yet answered Points out of 5 | The vertices of a regular nonagon (9-sided polygon) are to be labeled with the digits 1 through 9 in such a way that the sum of the numbers on every three consecutive vertices is a multiple of 3. Two acceptable arrangements are considered to be indistinguishable if one can be obtained from the other by rotating the nonagon in the plane. Find the number of distinguishable acceptable arrangements. |
|--|--|
| Question 6 Not yet answered Points out of 5 | Suppose that a parabola has vertex $(\frac{1}{4}, -\frac{9}{8})$ and equation $y = ax^2 + bx + c$, where $a > 0$ and $a + b + c$ is an integer. The minimum possible value of a can be written in the form $\frac{p}{q}$, where p and q are relatively prime positive integers. Find $p + q$. Answer: |
| Question 7 Not yet answered Points out of 5 | Find the number of positive integers m for which there exist nonnegative integers x_0, x_1 , , x_{2011} such that $m^{x_0} = \sum_{k=1}^{2011} m^{x_k}.$ Answer: |
| Question 8 Not yet answered Points out of 5 | In triangle ABC , $BC = 23$, $CA = 27$, and $AB = 30$. Points V and W are on \overline{AC} with V on \overline{AW} , points X and Y are on \overline{BC} with X on \overline{CY} , and points Z and U are on \overline{AB} with Z on \overline{BU} . In addition, the points are positioned so that $\overline{UV} \parallel \overline{BC}$, $\overline{WX} \parallel \overline{AB}$, and $\overline{YZ} \parallel \overline{CA}$. Right angle folds are then made along \overline{UV} , \overline{WX} , and \overline{YZ} . The resulting figure is placed on a level floor to make a table with triangular legs. Let h be the maximum possible height of a table constructed from triangle ABC whose top is parallel to the floor. Then h can be written in the form $\frac{k\sqrt{m}}{n}$, where k and n are relatively prime positive integers and m is a positive integer that is not divisible by the square of any prime. Find $k + m + n$. |

| Question 9 Not yet answered Points out of 5 | Suppose x is in the interval $[0, \pi/2]$ and $\log_{24 \sin x} (24 \cos x) = \frac{3}{2}$. Find $24 \cot^2 x$. Answer: |
|---|--|
| Question 10 Not yet answered Points out of 5 | The probability that a set of three distinct vertices chosen at random from among the vertices of a regular n-gon determine an obtuse triangle is $\frac{93}{125}$. Find the sum of all possible values of <i>n</i> . |
| Question 11 Not yet answered Points out of 5 | Let R be the set of all possible remainders when a number of the form 2^n , n a nonnegative integer, is divided by 1000. Let S be the sum of the elements in R . Find the remainder when S is divided by 1000. |
| Question 12 Not yet answered Points out of 5 | Six men and some number of women stand in a line in random order. Let p be the probability that a group of at least four men stand together in the line, given that every man stands next to at least one other man. Find the least number of women in the line such that p does not exceed 1 percent. |
| Question 13 Not yet answered Points out of 5 | A cube with side length 10 is suspended above a plane. The vertex closest to the plane is labeled A . The three vertices adjacent to vertex A are at heights 10, 11, and 12 above the plane. The distance from vertex A to the plane can be expressed as $\frac{r-\sqrt{s}}{t}$, where r , s , and t are positive integers, and $r + s + t < 1000$. Find $r + s + t$. |

| Question 14 Not yet answered Points out of 5 | Let $A_1A_2A_3A_4A_5A_6A_7A_8$ be a regular octagon. Let M_1 , M_3 , M_5 , and M_7 be the midpoints of sides $\overline{A_1A_2}$, $\overline{A_3A_4}$, $\overline{A_5A_6}$, and $\overline{A_7A_8}$, respectively. For $i = 1, 3, 5, 7$, ray R_i is constructed from M_i towards the interior of the octagon such that $R_1 \perp R_3$, $R_3 \perp R_5$, $R_5 \perp R_7$, and $R_7 \perp R_1$. Pairs of rays R_1 and R_3 , R_3 and R_5 , R_5 and R_7 , and $R_7 \perp R_1$. Pairs of rays R_1 and $R_3 = A_1A_2$, then $\cos 2\angle A_3M_3B_1$ can be written in the form $m - \sqrt{n}$, where m and n are positive integers. Find $m + n$. |
|---|--|
| | |
| Question 15 Not yet answered | For some integer m , the polynomial $x^3-2011x+m$ has the three integer roots $a,b,$ and $c.$ Find $ a + b + c .$ |
| Points out of 5 | Answer: |