

## 2012 AIME I

For more practice and resources, visit ziml.areteem.org

The problems in the AMC-Series Contests are copyrighted by American Mathematics Competitions at Mathematical Association of America (www.maa.org).



Question <b>1</b> Not yet answered Points out of 5	Find the number of positive integers with three not necessarily distinct digits, $abc$ , with $a \neq 0$ and $c \neq 0$ such that both $abc$ and $cba$ are multiples of 4.
Question 2 Not yet answered Points out of 5	The terms of an arithmetic sequence add to 715. The first term of the sequence is increased by 1, the second term is increased by 3, the third term is increased by 5, and in general, the $k$ th term is increased by the $k$ th odd positive integer. The terms of the new sequence add to 836. Find the sum of the first, last, and middle terms of the original sequence.
<b>Question 3</b> Not yet answered Points out of 5	Nine people sit down for dinner where there are three choices of meals. Three people order the beef meal, three order the chicken meal, and three order the fish meal. The waiter serves the nine meals in random order. Find the number of ways in which the waiter could serve the meal types to the nine people so that exactly one person receives the type of meal ordered by that person.
Question 4 Not yet answered Points out of 5	Butch and Sundance need to get out of Dodge. To travel as quickly as possible, each alternates walking and riding their only horse, Sparky, as follows. Butch begins by walking while Sundance rides. When Sundance reaches the first of the hitching posts that are conveniently located at one-mile intervals along their route, he ties Sparky to the post and begins walking. When Butch reaches Sparky, he rides until he passes Sundance, then leaves Sparky at the next hitching post and resumes walking, and they continue in this manner. Sparky, Butch, and Sundance walk at 6, 4, and 2.5 miles per hour, respectively. The first time Butch and Sundance meet at a milepost, they are $n$ miles from Dodge, and they have been traveling for $t$ minutes. Find $n + t$ .

Question 5 Let B be the set of all binary integers that can be written using exactly 5 zeros and 8 ones where leading zeros are allowed. If all possible subtractions are performed in which one Not yet answered element of B is subtracted from another, find the number of times the answer 1 is obtained. Points out of 5 Answer: Question 6 The complex numbers z and w satisfy  $z^{13} = w, w^{11} = z$ , and the imaginary part of z is  $\sin rac{m\pi}{n}$  , for relatively prime positive integers m and n with m < n. Find n.Not yet answered Points out of 5 Answer: Question 7 At each of the sixteen circles in the network below stands a student. A total of 3360 coins are distributed among the sixteen students. All at once, all students give away all their coins Not yet answered by passing an equal number of coins to each of their neighbors in the network. After the Points out of 5 trade, all students have the same number of coins as they started with. Find the number of coins the student standing at the center circle had originally. Answer:





Question 9 Not yet answered Points out of 5	Let $x, y,$ and $z$ be positive real numbers that satisfy
	$2\log_x(2y)=2\log_{2x}(4z)=\log_{2x^4}(8yz) eq 0.$
	The value of $xy^5z$ can be expressed in the form $rac{1}{2^{p/q}},$ where $p$ and $q$ are relatively prime
	positive integers. Find $p+q.$
	Answer:

Question <b>10</b> Not yet answered Points out of 5	Let $S$ be the set of all perfect squares whose rightmost three digits in base 10 are 256. Let $\mathcal{T}$ be the set of all numbers of the form $\frac{x-256}{1000}$ , where $x$ is in $S$ . In other words, $\mathcal{T}$ is the set of numbers that result when the last three digits of each number in $S$ are truncated. Find the remainder when the tenth smallest element of $\mathcal{T}$ is divided by 1000.
Question <b>11</b> Not yet answered Points out of 5	A frog begins at $P_0 = (0, 0)$ and makes a sequence of jumps according to the following rule: from $P_n = (x_n, y_n)$ , the frog jumps to $P_{n+1}$ , which may be any of the points $(x_n + 7, y_n + 2), (x_n + 2, y_n + 7), (x_n - 5, y_n - 10),$ or $(x_n - 10, y_n - 5)$ . There are $M$ points $(x, y)$ with $ x  +  y  \le 100$ that can be reached by a sequence of such jumps. Find the remainder when $M$ is divided by 1000.
Question 12 Not yet answered Points out of 5	Let $\triangle ABC$ be a right triangle with right angle at $C$ . Let $D$ and $E$ be points on $\overline{AB}$ with $D$ between $A$ and $E$ such that $\overline{CD}$ and $\overline{CE}$ trisect $\angle C$ . If $\frac{DE}{BE} = \frac{8}{15}$ , then $\tan B$ can be written as $\frac{m\sqrt{p}}{n}$ , where $m$ and $n$ are relatively prime positive integers, and $p$ is a positive integer not divisible by the square of any prime. Find $m + n + p$ . Answer:
Question 13 Not yet answered Points out of 5	Three concentric circles have radii 3, 4, and 5. An equilateral triangle with one vertex on each circle has side length <i>s</i> . The largest possible area of the triangle can be written as $a + \frac{b}{c}\sqrt{d}$ , where <i>a</i> , <i>b</i> , <i>c</i> , and <i>d</i> are positive integers, <i>b</i> and <i>c</i> are relatively prime, and <i>d</i> is not divisible by the square of any prime. Find $a + b + c + d$ . Answer:
Question 14 Not yet answered Points out of 5	Complex numbers $a$ , $b$ , and $c$ are zeros of a polynomial $P(z) = z^3 + qz + r$ , and $ a ^2 +  b ^2 +  c ^2 = 250$ . The points corresponding to $a$ , $b$ , and $c$ in the complex plane are the vertices of a right triangle with hypotenuse $h$ . Find $h^2$ .

<b>Question 15</b> Not yet answered Points out of 5	There are $n$ mathematicians seated around a circular table with $n$ seats numbered $1, 2, 3, \ldots, n$ in clockwise order. After a break they again sit around the table. The mathematicians note that there is a positive integer $a$ such that:
	(1) for each $k$ , the mathematician who was seated in seat $k$ before the break is seated in seat $ka$ after the break (where seat $i+n$ is seat $i$ ), and
	(2) for every pair of mathematicians, the number of mathematicians sitting between them after the break, counting in both the clockwise and the counterclockwise directions, is different from either of the number of mathematicians sitting between them before the break.
	Find the number of possible values of $n$ with $1 < n < 1000.$
	Answer: