



2012 AIME II

For more practice and resources, visit ziml.aretteam.org

The problems in the AMC-Series Contests are copyrighted
by American Mathematics Competitions at Mathematical
Association of America (www.maa.org).



Question 1

Not yet answered

Points out of 5

Find the number of ordered pairs of positive integer solutions (m, n) to the equation $20m + 12n = 2012$.

Answer: **Question 2**

Not yet answered

Points out of 5

Two geometric sequences a_1, a_2, a_3, \dots and b_1, b_2, b_3, \dots have the same common ratio, with $a_1 = 27$, $b_1 = 99$, and $a_{15} = b_{11}$. Find a_9 .

Answer: **Question 3**

Not yet answered

Points out of 5

At a certain university, the division of mathematical sciences consists of the departments of mathematics, statistics, and computer science. There are two male and two female professors in each department. A committee of six professors is to contain three men and three women and must also contain two professors from each of the three departments. Find the number of possible committees that can be formed subject to these requirements.

Answer: **Question 4**

Not yet answered

Points out of 5

Ana, Bob, and Cao bike at constant rates of 8.6 meters per second, 6.2 meters per second, and 5 meters per second, respectively. They all begin biking at the same time from the northeast corner of a rectangular field whose longer side runs due west. Ana starts biking along the edge of the field, initially heading west, Bob starts biking along the edge of the field, initially heading south, and Cao bikes in a straight line across the field to a point D on the south edge of the field. Cao arrives at point D at the same time that Ana and Bob arrive at D for the first time. The ratio of the field's length to the field's width to the distance from point D to the southeast corner of the field can be represented as $p : q : r$, where p , q , and r are positive integers with p and q relatively prime. Find $p + q + r$.


Answer:

Question 5

Not yet answered

Points out of 5

In the accompanying figure, the outer square S has side length 40. A second square S' of side length 15 is constructed inside S with the same center as S and with sides parallel to those of S . From each midpoint of a side of S , segments are drawn to the two closest vertices of S' . The result is a four-pointed starlike figure inscribed in S . The star figure is cut out and then folded to form a pyramid with base S' .



Find the volume of this pyramid.

Answer:

Question 6

Not yet answered

Points out of 5

Let $z = a + bi$ be the complex number with $|z| = 5$ and $b > 0$ such that the distance between $(1 + 2i)z^3$ and z^5 is maximized, and let $z^4 = c + di$. Find $c + d$.

Answer:

Question 7

Not yet answered

Points out of 5

Let S be the increasing sequence of positive integers whose binary representation has exactly 8 ones. Let N be the 1000th number in S . Find the remainder when N is divided by 1000.

Answer:

Question 8

Not yet answered

Points out of 5

The complex numbers z and w satisfy the system

$$z + \frac{20i}{w} = 5 + i$$

$$w + \frac{12i}{z} = -4 + 10i$$

Find the smallest possible value of $|zw|^2$.

Answer:**Question 9**

Not yet answered

Points out of 5

Let x and y be real numbers such that $\frac{\sin x}{\sin y} = 3$ and $\frac{\cos x}{\cos y} = \frac{1}{2}$. The value of $\frac{\sin 2x}{\sin 2y} + \frac{\cos 2x}{\cos 2y}$ can be expressed in the form $\frac{p}{q}$, where p and q are relatively prime positive integers. Find $p + q$.

Answer:**Question 10**

Not yet answered

Points out of 5

Find the number of positive integers n less than 1000 for which there exists a positive real number x such that $n = x \lfloor x \rfloor$.

Note: $\lfloor x \rfloor$ is the greatest integer less than or equal to x .

Answer:**Question 11**

Not yet answered

Points out of 5

Let $f_1(x) = \frac{2}{3} - \frac{3}{3x+1}$, and for $n \geq 2$, define $f_n(x) = f_1(f_{n-1}(x))$. The value of x that satisfies $f_{1001}(x) = x - 3$ can be expressed in the form $\frac{m}{n}$, where m and n are relatively prime positive integers. Find $m + n$.

Answer:

Question 12

Not yet answered

Points out of 5

For a positive integer p , define the positive integer n to be p -safe if n differs in absolute value by more than 2 from all multiples of p . For example, the set of 10-safe numbers is $\{3, 4, 5, 6, 7, 13, 14, 15, 16, 17, 23, \dots\}$. Find the number of positive integers less than or equal to 10,000 which are simultaneously 7-safe, 11-safe, and 13-safe.

Answer:

Question 13

Not yet answered

Points out of 5

Equilateral $\triangle ABC$ has side length $\sqrt{111}$. There are four distinct triangles AD_1E_1 , AD_1E_2 , AD_2E_3 , and AD_2E_4 , each congruent to $\triangle ABC$, with $BD_1 = BD_2 = \sqrt{11}$. Find $\sum_{k=1}^4 (CE_k)^2$.

Answer:

Question 14

Not yet answered

Points out of 5

In a group of nine people each person shakes hands with exactly two of the other people from the group. Let N be the number of ways this handshaking can occur. Consider two handshaking arrangements different if and only if at least two people who shake hands under one arrangement do not shake hands under the other arrangement. Find the remainder when N is divided by 1000.

Answer:

Question 15

Not yet answered

Points out of 5

Triangle ABC is inscribed in circle ω with $AB = 5$, $BC = 7$, and $AC = 3$. The bisector of angle A meets side \overline{BC} at D and circle ω at a second point E . Let γ be the circle with diameter \overline{DE} . Circles ω and γ meet at E and a second point F . Then $AF^2 = \frac{m}{n}$, where m and n are relatively prime positive integers. Find $m + n$.

Answer: