

2021 AIME II

For more practice and resources, visit ziml.areteem.org

The problems in the AMC-Series Contests are copyrighted by American Mathematics Competitions at Mathematical Association of America (www.maa.org).



Question 1 Not yet answered Points out of 1	Find the arithmetic mean of all the three-digit palindromes. (Recall that a palindrome is a number that reads the same forward and backward, such as 777 or 383.) Answer:
Question 2 Not yet answered Points out of 1	Equilateral triangle ABC has side length 840. Point D lies on the same side of line BC as A such that $\overline{BD} \perp \overline{BC}$. The line ℓ through D parallel to line BC intersects sides \overline{AB} and \overline{AC} at points E and F , respectively. Point G lies on ℓ such that F is between E and G , $\triangle AFG$ is isosceles, and the ratio of the area of $\triangle AFG$ to the area of $\triangle BED$ is $8:9$. Find AF . Answer:
Question 3 Not yet answered Points out of 1	Find the number of permutations x_1, x_2, x_3, x_4, x_5 of numbers $1, 2, 3, 4, 5$ such that the sum of five products $x_1x_2x_3 + x_2x_3x_4 + x_3x_4x_5 + x_4x_5x_1 + x_5x_1x_2$ is divisible by 3. Answer:
Question 4 Not yet answered Points out of 1	There are real numbers a , b , c , and d such that -20 is a root of $x^3 + ax + b$ and -21 is a root of $x^3 + cx^2 + d$. These two polynomials share a complex root $m + \sqrt{n} \cdot i$, where m and n are positive integers and $i = \sqrt{-1}$. Find $m + n$.
Question 5 Not yet answered Points out of 1	For positive real numbers s , let $\tau(s)$ denote the set of all obtuse triangles that have area s and two sides with lengths 4 and 10. The set of all s for which $\tau(s)$ is nonempty, but all triangles in $\tau(s)$ are congruent, is an interval $[a, b)$. Find $a^2 + b^2$.
Question 6 Not yet answered Points out of 1	For any finite set S , let $ S $ denote the number of elements in S . Find the number of ordered pairs (A, B) such that A and B are (not necessarily distinct) subsets of $\{1, 2, 3, 4, 5\}$ that satisfy $ A \cdot B = A \cap B \cdot A \cup B .$ Answer:

Question 7		
Not yet answered		

Points out of 1

Let *a*, *b*, *c*, and *d* be real numbers that satisfy the system of equations

 $a+b=-3,\ ab+bc+ca=-4,\ abc+bcd+cda+dab=14,\ abcd=30.$

There exist relatively prime positive integers m and n such that

$$a^2+b^2+c^2+d^2=rac{m}{n}.$$

Find m + n.

Answer:

Question 8

Not yet answered

Points out of 1

An ant makes a sequence of moves on a cube where a move consists of walking from one vertex to an adjacent vertex along an edge of the cube. Initially the ant is at a vertex of the bottom face of the cube and chooses one of the three adjacent vertices to move to as its first move. For all moves after the first move, the ant does not return to its previous vertex, but chooses to move to one of the other two adjacent vertices. All choices are selected at random so that each of the possible moves is equally likely. The probability that after exactly 8 moves that ant is at a vertex of the top face on the cube is $\frac{m}{n}$, where m and n are relatively prime positive integers. Find m + n.

Answer:

Question 9	Find the number of ordered pairs (m, n) such that m and n are positive integers in the set $\{1, 2,, 30\}$
Not yet answered	and the greatest common divisor of $2^m + 1$ and $2^n - 1$ is not 1.
Points out of 1	Answer:
Question 10 Not yet answered Points out of 1	Two spheres with radii 36 and one sphere with radius 13 are each externally tangent to the other two spheres and to two different planes \mathcal{P} and \mathcal{Q} . The intersection of planes \mathcal{P} and \mathcal{Q} is the line ℓ . The distance from line ℓ to the point where the sphere with radius 13 is tangent to plane \mathcal{P} is $\frac{m}{n}$, where m and n are relatively prime positive integers. Find $m + n$.

ut of 1 the

Question 11 Not yet answered Points out of 1	A teacher was leading a class of four perfectly logical students. The teacher chose a set S of four integers and gave a different number in S to each student. Then the teacher announced to the class that the numbers in S were four consecutive two-digit positive integers, that some number in S was divisible by 6, and a different number in S was divisible by 7. The teacher then asked if any of the students could deduce what S is, but in unison, all of the students replied no. However, upon hearing that all four students replied no, each student was able to determine the elements of S. Find the sum of all possible values of the greatest element of S .
Question 12 Not yet answered Points out of 1	A convex quadrilateral has area 30 and side lengths 5, 6, 9, and 7, in that order. Denote by θ the measure of the acute angle formed by the diagonals of the quadrilateral. Then $\tan \theta$ can be written in the form $\frac{m}{n}$, where m and n are relatively prime positive integers. Find $m + n$.
Question 13 Not yet answered Points out of 1	Find the least positive integer n for which $2^n + 5^n - n$ is a multiple of 1000. Answer:
Question 14 Not yet answered Points out of 1	Let ΔABC be an acute triangle with circumcenter O and centroid G . Let X be the intersection of the line tangent to the circumcircle of ΔABC at A and the line perpendicular to GO at G . Let Y be the intersection of lines XG and BC . Given that the measures of $\angle ABC$, $\angle BCA$, and $\angle XOY$ are in the ratio $13:2:17$, the degree measure of $\angle BAC$ can be written as $\frac{m}{n}$, where m and n are relatively prime positive integers. Find $m + n$.
Question 15 Not yet answered Points out of 1	Let $f(n)$ and $g(n)$ be functions satisfying $f(n) = \left\{ egin{array}{cc} \sqrt{n}, & ext{if } \sqrt{n} ext{ is an integer}, \ 1+f(n+1), & ext{otherwise}. \end{array} ight.$
	and
	$g(n) = egin{cases} \sqrt{n}, & ext{if } \sqrt{n} ext{ is an integer}, \ 2+g(n+2), & ext{otherwise}. \end{cases}$
	for positive integers n. Find the least positive integer n such that $\frac{f(n)}{g(n)} = \frac{4}{7}$.
	Answer: